

1. In the following reaction
- $$\text{NH}_4^+(\text{aq}) + \text{H}_2\text{O}(\text{l}) \leftrightarrow \text{NH}_3(\text{aq}) + \text{H}_3\text{O}^+(\text{aq})$$
- $\text{NH}_4^+$  is an acid and  $\text{NH}_3$  is its conjugate base.
  - $\text{H}_2\text{O}$  is an acid and  $\text{H}_3\text{O}^+$  is its conjugate base.
  - $\text{NH}_4^+$  is an acid and  $\text{H}_3\text{O}^+$  is its conjugate base.
  - $\text{H}_2\text{O}$  is an acid and  $\text{NH}_4^+$  is its conjugate base.
  - $\text{NH}_3$  is an acid and  $\text{NH}_4^+$  is its conjugate base.
2. Which of the following acids has the strongest conjugate base?
- Ascorbic acid,  $K_a = 8.0 \times 10^{-5}$
  - Benzoic acid,  $K_a = 6.5 \times 10^{-5}$
  - 3-chlorobenzoic acid,  $K_a = 1.5 \times 10^{-4}$
  - 2-hydroxybenzoic acid,  $K_a = 1.1 \times 10^{-3}$
  - Chloroacetic acid,  $K_a = 1.4 \times 10^{-3}$
3. Knowing that  $\text{H}_2\text{S}$  is a stronger acid than  $\text{HCN}$ , determine, if possible, in which direction the following equilibrium lies.
- $$\text{HCN}(\text{aq}) + \text{HS}^-(\text{aq}) \leftrightarrow \text{CN}^-(\text{aq}) + \text{H}_2\text{S}(\text{aq})$$
- equilibrium lies to the left
  - equilibrium lies to the right
  - equilibrium is perfectly balanced left and right
  - can be determined if the relative acidity of  $\text{HS}^-$  is given
  - cannot be determined
4. What is the pH of a 0.054 M NaOH solution at 25 °C?
- 1.14
  - 1.27
  - 8.64
  - 12.73
  - 13.95
5. We dilute 1.00 mL of 1.00 M HCl solution to 100.0 mL. What is  $[\text{OH}^-]$  in this solution at 25 °C?
- $1.00 \times 10^{12}$  M
  - $1 \times 10^2$  M
  - 0.010 M
  - $7.00 \times 10^{-4}$  M
  - $1.00 \times 10^{-12}$  M
6. At 25 °C, what is the pH of a 1.75 M solution of sodium cyanide NaCN? ( $K_b = 2.5 \times 10^{-5}$ )
- 11.82
  - 10.04
  - 3.44
  - 2.18
  - 0.80
7. At 25 °C, what is the pH of a 3.25 M solution of ammonium chloride,  $\text{NH}_4\text{Cl}$ ? ( $K_a = 5.6 \cdot 10^{-10}$ )
- 2.37
  - 4.37
  - 4.62
  - 9.37
  - 9.63
8. Which of the following acid-base reactions will lie predominantly toward the products? ( $K_a(\text{CH}_3\text{CO}_2\text{H}) = 1.8 \cdot 10^{-5}$ ,  $K_b(\text{NH}_3) = 1.8 \cdot 10^{-5}$ )
- Reaction 1:  $\text{NH}_3(\text{aq}) + \text{H}_2\text{O}(\text{l}) \leftrightarrow \text{NH}_4^+(\text{aq}) + \text{OH}^-(\text{aq})$
- Reaction 2:  $\text{CH}_3\text{CO}_2\text{H}(\text{aq}) + \text{H}_2\text{O}(\text{l}) \leftrightarrow \text{H}_3\text{O}^+(\text{aq}) + \text{CH}_3\text{CO}_2^-(\text{aq})$
- Reaction 3:  $\text{CH}_3\text{CO}_2\text{H}(\text{aq}) + \text{NH}_3(\text{aq}) \leftrightarrow \text{NH}_4^+(\text{aq}) + \text{CH}_3\text{CO}_2^-(\text{aq})$
- 1 only
  - 2 only
  - 3 only
  - 1 and 2 only
  - 1 and 3 only


9. If you mix 250. mL of 0.24 M HF with 75.0 mL of 0.80 M NaOH, what is the pH of the resulting solution? For  $F^-$ ,  $K_b = 1.4 \times 10^{-11}$
- 5.42
  - 5.79
  - 6.24
  - 7.53
  - 8.21
10. If you mix 125. mL of 0.50 M  $CH_3CO_2H$  with 75.0 mL of 0.83 M NaOH, what is the pH of the resulting solution? For  $CH_3COOH$ ,  $K_a = 1.8 \times 10^{-5}$
- 4.88
  - 5.01
  - 7.14
  - 8.99
  - 9.76
11. If you mix equal molar quantities of NaOH and  $CH_3CO_2H$ , what are the principal species present in the resulting solution?
- $Na^+$ ,  $CH_3CO_2^-$ ,  $OH^-$ , and  $H_2O$
  - $Na^+$ ,  $CH_3CO_2^-$ ,  $CH_3CO_2H$ ,  $OH^-$ , and  $H_2O$
  - $Na^+$ ,  $CH_3CO_2H$ ,  $OH^-$ , and  $H_2O$
  - $Na^+$ ,  $CH_3CO_2^-$ ,  $H_3O^+$ , and  $H_2O$
  - $Na^+$ ,  $CH_3CO_2H$ ,  $H_3O^+$ , and  $H_2O$
12. The salt produced by the reaction of an equal number of moles of KOH and  $HNO_3$  will react with water to give a solution which is
- acidic.
  - basic.
  - neutral.
  - non-ionic.
  - impossible to determine.
13. If you mix equal molar quantities of  $NH_3$  ( $K_b = 1.8 \times 10^{-5}$ ) and  $CH_3CO_2H$  ( $K_a = 1.8 \times 10^{-5}$ ), the resulting solution will be
- acidic because  $K_a$  of  $NH_4^+$  is greater than  $K_b$  of  $CH_3CO_2^-$ .
  - acidic because  $K_a$  of  $NH_4^+$  is greater than  $K_a$  of  $CH_3CO_2H$ .
  - basic because  $K_b$  of  $NH_3$  is greater than  $K_b$  of  $CH_3CO_2^-$ .
  - basic because  $K_a$  of  $NH_4^+$  is greater than  $K_b$  of  $CH_3CO_2^-$ .
  - neutral because  $K_a$  of  $NH_4^+$  equals  $K_b$  of  $CH_3CO_2^-$ .
14. At the neutralization point of the titration of an acid with base, what condition is met?
- Volume of base added from buret equals volume acid in reaction flask.
  - Molarity of base from the buret equals molarity of acid in reaction flask.
  - Moles of base added from the buret equals moles of acid in the reaction flask.
  - % ionization of base added from the buret equals % ionization of the acid in flask.
  - All of the above conditions are met.
15. The solubility of  $FeCO_3$  is  $5.9 \times 10^{-6}$  mol/L. What is  $K_{sp}$  for  $FeCO_3$ ?
- $5.9 \times 10^{-6}$
  - $1.2 \times 10^{-21}$
  - $3.5 \times 10^{-11}$
  - $2.8 \times 10^{-10}$
  - $1.3 \times 10^{-14}$

16. What is the concentration of  $\text{CrO}_4^{2-}$  in a saturated solution of  $\text{PbCrO}_4$  if  $K_{\text{sp}} = 1.8 \times 10^{-14}$ ?
- $1.3 \times 10^{-7} \text{ M}$
  - $7.5 \times 10^{-6} \text{ M}$
  - $1.8 \times 10^{-4} \text{ M}$
  - $1.3 \times 10^{-4} \text{ M}$
  - $5.1 \times 10^{-3} \text{ M}$
17. Which of the following has the highest molar solubility?
- $\text{PbCO}_3$ ;  $K_{\text{sp}} = 1.5 \times 10^{-13}$
  - $\text{PbS}$ ;  $K_{\text{sp}} = 8.4 \times 10^{-28}$
  - $\text{PbI}_2$ ;  $K_{\text{sp}} = 8.7 \times 10^{-9}$
  - $\text{PbSO}_4$ ;  $K_{\text{sp}} = 1.8 \times 10^{-8}$
  - $\text{Pb}_2(\text{PO}_4)_2$ ;  $K_{\text{sp}} = 3.0 \times 10^{-44}$
18. Calculate the equilibrium constant for the reaction:
- $$\text{CdS(s)} + \text{Zn}^{2+}(\text{aq}) \leftrightarrow \text{ZnS(s)} + \text{Cd}^{2+}(\text{aq})$$
- $\text{CdS}$ ;  $K_{\text{sp}} = 3.6 \times 10^{-29}$     $\text{ZnS}$ ;  $K_{\text{sp}} = 1.1 \times 10^{-21}$
- $3.3 \times 10^{-8}$
  - $2.7 \times 10^{-4}$
  - $4.2 \times 10^5$
  - $2.5 \times 10^{49}$
  - $3.1 \times 10^7$
19. For  $\text{Ag}_2\text{SO}_4$ ,  $K_{\text{sp}} = 1.7 \times 10^{-5}$ . How many grams of  $\text{Na}_2\text{SO}_4$  (MM = 142.0 g/mol) must be added to 100. mL of 0.022 M  $\text{AgNO}_3$  to just initiate precipitation?
- 5.0 g
  - 4.0 g
  - 3.0 g
  - 0.50 g
  - 0.40 g
20. For thallium bromide,  $\text{TlBr}$ ,  $K_{\text{sp}} = 3.4 \times 10^{-6}$ . How many grams of  $\text{KBr}$  (MM = 119.0 g/mol) must be added to 100. mL of  $5.5 \times 10^{-4} \text{ M TlNO}_3$  to just initiate precipitation?
- 0.74 g
  - 0.074 g
  - 0.065 g
  - 0.0065 g
  - 0.0033 g
21. In the following reaction
- $$\text{HF(aq)} + \text{H}_2\text{O(l)} \leftrightarrow \text{H}_3\text{O}^+(\text{aq}) + \text{F}^-(\text{aq})$$
- $\text{HF}$  is an acid and  $\text{H}_3\text{O}^+$  is its conjugate base.
  - $\text{H}_2\text{O}$  is an acid and  $\text{H}_3\text{O}^+$  is its conjugate base.
  - $\text{HF}$  is an acid and  $\text{F}^-$  is its conjugate base.
  - $\text{H}_2\text{O}$  is an acid and  $\text{H}_3\text{O}^+$  is its conjugate base.
  - $\text{HF}$  is an acid and  $\text{H}_2\text{O}$  is its conjugate base.
22. What is the pH of a  $4.2 \times 10^{-4} \text{ M HBr}$  solution at  $25^\circ\text{C}$ ?
- 2.80
  - 3.38
  - 3.80
  - 4.20
  - 4.62
23. We have a  $4.63 \times 10^{-4} \text{ M}$  solution of  $\text{HCl}$ . What is the pH of this solution at  $25^\circ\text{C}$ ?
- 3.33
  - 4.00
  - 4.63
  - 8.37
  - 9.25

25. What is the pH of a 3.18 M  $\text{CH}_3\text{COOH}$  solution at 25 °C?  $K_a = 1.8 \times 10^{-5}$ ?
- 2.12
  - 2.75
  - 1.40
  - 4.24
  - 4.74
26. What is the % ionization of a 3.14 M  $\text{CH}_3\text{CO}_2\text{H}$  solution at 25 °C? For  $\text{CH}_3\text{CO}_2\text{H}$ ,  $K_a = 1.8 \times 10^{-5}$ .
- 0.24%
  - 0.57%
  - 1.8%
  - 3.2%
  - 7.5%
27. Which of the following acid-base reactions will lie predominantly toward the products?
- Reaction 1:  $\text{HF}(\text{aq}) + \text{NH}_3(\text{aq}) \leftrightarrow \text{NH}_4^+(\text{aq}) + \text{F}^-(\text{aq})$
- Reaction 2:  $\text{NH}_3(\text{aq}) + \text{H}_2\text{O}(\text{l}) \leftrightarrow \text{NH}_4^+(\text{aq}) + \text{OH}^-(\text{aq})$
- Reaction 3:  $\text{HF}(\text{aq}) + \text{H}_2\text{O}(\text{l}) \leftrightarrow \text{H}_3\text{O}^+(\text{aq}) + \text{F}^-(\text{aq})$
- 1 only
  - 2 only
  - 1 and 2 only
  - 2 and 3 only
  - 1, 2, and 3
28. We add 1.00 mL of 10.0 M  $\text{HNO}_3$  to 100. mL of 0.10 M  $\text{NaHCOO}$ . What is the pH of the resulting solution?  $K_a(\text{HCOOH}) = 1.8 \times 10^{-4}$
- 2.37
  - 3.45
  - 4.27
  - 4.35
  - 11.60
29. If you mix 100. mL of 0.11 M  $\text{HCl}$  with 50.0 mL of 0.22 M  $\text{NH}_3$ , what is the pH of the resulting solution? For  $\text{NH}_4^+$ ,  $K_a = 5.6 \times 10^{-10}$
- 4.63
  - 5.19
  - 6.02
  - 8.37
  - 9.37
30. If you mix 125. mL of 0.50 M  $\text{CH}_3\text{CO}_2\text{H}$  with 75.0 mL of 0.83 M  $\text{NaOH}$ , what is the pH of the resulting solution? For  $\text{CH}_3\text{COO}^-$ ,  $K_b = 5.6 \times 10^{-10}$
- 4.88
  - 5.01
  - 7.14
  - 9.12
  - 9.76
31. What effect will the addition of the reagent in each of the following have on the pH of the  $\text{CH}_3\text{CO}_2\text{H}$  solution respectively?
- Flask 1: Addition of  $\text{NaCH}_3\text{CO}_2$  to  $\text{CH}_3\text{CO}_2\text{H}(\text{aq})$
- Flask 2: Addition of  $\text{Ca}(\text{CH}_3\text{CO}_2)_2$  to  $\text{CH}_3\text{CO}_2\text{H}(\text{aq})$
- no change, increase
  - no change, decrease
  - decrease, no change
  - decrease, decrease
  - increase, increase
32. If you add 20.0 mL of 2.30 M  $\text{NH}_3$  to 100. mL of a 1.17 M  $\text{NH}_4\text{Cl}$  solution, what is the pH of the resulting solution? For  $\text{NH}_3$ ,  $K_b = 1.8 \times 10^{-5}$
- 5.15
  - 6.35
  - 7.10
  - 7.65
  - 8.85

33. We have 250. mL of a 0.56 M solution of  $\text{NaCH}_3\text{COO}$ . How many milliliters of a 0.50 M  $\text{CH}_3\text{COOH}$  solution should be added to make a buffer of  $\text{pH} = 4.40$ ?  
 $K_a(\text{CH}_3\text{COOH}) = 1.8 \times 10^{-5}$
- 200
  - 230
  - 620
  - 710
  - 750
34. Which of the following is the solubility product constant for  $\text{Mn}(\text{OH})_2$ ?
- $K_{\text{sp}} = [\text{Mn}^{2+}][\text{OH}^-]^2$
  - $K_{\text{sp}} = [\text{Mn}^{2+}][2\text{OH}^-]^2$
  - $K_{\text{sp}} = [\text{Mn}^{2+}]^2[\text{OH}^-]^2$
  - $K_{\text{sp}} = [\text{Mn}^{2+}]^2[\text{OH}^-]$
  - $K_{\text{sp}} = [\text{Mn}^{2+}]^2[\text{OH}^-]^2$
35. Rank the compounds from lowest to highest molar solubility.  
 $\text{FeCO}_3$ ;  $K_{\text{sp}} = 3.5 \times 10^{-11}$   
 $\text{BaSO}_4$ ;  $K_{\text{sp}} = 1.1 \times 10^{-10}$   
 $\text{ZnCO}_3$ ;  $K_{\text{sp}} = 1.5 \times 10^{-11}$
- $\text{ZnCO}_3 < \text{BaSO}_4 < \text{FeCO}_3$
  - $\text{FeCO}_3 < \text{ZnCO}_3 < \text{BaSO}_4$
  - $\text{ZnCO}_3 < \text{FeCO}_3 < \text{BaSO}_4$
  - $\text{BaSO}_4 < \text{ZnCO}_3 < \text{FeCO}_3$
  - $\text{BaSO}_4 < \text{FeCO}_3 < \text{ZnCO}_3$
36. What is the concentration of  $\text{SO}_4^{2-}$  in a saturated solution of  $\text{BaSO}_4$  if  $K_{\text{sp}} = 1.1 \times 10^{-10}$ ?
- $1.1 \times 10^{-10}$  M
  - $5.5 \times 10^{-11}$  M
  - $5.0 \times 10^{-5}$  M
  - $1.0 \times 10^{-5}$  M
  - $9.5 \times 10^{-4}$  M
37. For  $\text{MgF}_2$ ,  $K_{\text{sp}} = 6.4 \times 10^{-9}$ . If you mix 400. mL of  $1 \times 10^{-4}$  M  $\text{Mg}(\text{NO}_3)_2$  and 500. mL of  $1.00 \times 10^{-4}$  M  $\text{NaF}$ , what will be observed?
- A precipitate forms because  $Q_{\text{sp}} > K_{\text{sp}}$ .
  - A precipitate forms because  $Q_{\text{sp}} < K_{\text{sp}}$ .
  - No precipitate forms because  $Q_{\text{sp}} = K_{\text{sp}}$ .
  - No precipitate forms because  $Q_{\text{sp}} < K_{\text{sp}}$ .
  - No precipitate forms because  $Q_{\text{sp}} > K_{\text{sp}}$ .
38. For  $\text{AgI}$ ,  $K_{\text{sp}} = 8.3 \times 10^{-17}$ . What is the molar solubility of  $\text{AgI}$  in a solution which is  $5.1 \times 10^{-4}$  M in  $\text{AgNO}_3$ ?
- $5.1 \times 10^{-2}$  mol/L
  - $1.1 \times 10^{-5}$  mol/L
  - $8.3 \times 10^{-11}$  mol/L
  - $1.6 \times 10^{-13}$  mol/L
  - $4.2 \times 10^{-20}$  mol/L
39. Which of the following represents an increase in entropy?
- freezing of water
  - boiling of water
  - crystallization of salt from a supersaturated solution
  - the reaction  $2\text{NO}(\text{g}) \rightarrow \text{N}_2\text{O}_2(\text{g})$
  - the reaction  $2\text{H}_2(\text{g}) + \text{O}_2(\text{g}) \rightarrow 2\text{H}_2\text{O}(\text{g})$
40. If  $\Delta H$  and  $\Delta S$  are both negative or positive, then  $\Delta G$  has a \_\_\_\_\_ sign.
- positive
  - negative
  - variable
  - large
  - no


41. Calculate the standard entropy change for the following reaction,



given that  $S^\circ[\text{Cu(s)}] = 33.15$   
 $\text{J/K}\cdot\text{mol}$ ,  $S^\circ[\text{O}_2(\text{g})] = 205.14$   
 $\text{J/K}\cdot\text{mol}$ , and  $S^\circ[\text{CuO(s)}] = 42.63$   
 $\text{J/K}\cdot\text{mol}$

- 195.66 J/K
- 93.09 J/K
- 45.28 J/K
- 93.09 J/K
- 195.66 J/K

42. Calculate the standard entropy change for the following reaction,



given that  $S^\circ[\text{CCl}_4(\text{l})] = 216.40$   
 $\text{J/K}\cdot\text{mol}$ ,  $S^\circ[\text{CO}_2(\text{g})] = 213.74$   
 $\text{J/K}\cdot\text{mol}$ ,  $S^\circ[\text{O}_2(\text{g})] = 205.14$   
 $\text{J/K}\cdot\text{mol}$ , and  $S^\circ[\text{Cl}_2(\text{g})] = 223.07$   
 $\text{J/K}\cdot\text{mol}$ .

- 25.78 J/K
- 15.27 J/K
- +1.93 J/K
- 238.34 J/K
- 317.42 J/K

43. In which of the following reactions do you expect to have the smallest entropy change?

- $2\text{HF(g)} + \text{Cl}_2(\text{g}) \rightarrow 2\text{HCl(g)} + \text{F}_2(\text{g})$
- $2\text{Fe(s)} + 3/2 \text{O}_2(\text{g}) \rightarrow \text{Fe}_2\text{O}_3(\text{s})$
- $\text{CH}_4(\text{g}) + 2 \text{O}_2(\text{g}) \rightarrow \text{CO}_2(\text{g}) + 2\text{H}_2\text{O(l)}$
- $\text{Cu(s)} + 1/2 \text{O}_2(\text{g}) \rightarrow \text{CuO(s)}$
- $\text{H}_2(\text{g}) + \text{I}_2(\text{g}) \rightarrow 2\text{HI(g)}$


44. If  $\Delta G$  is positive at all temperatures, then  $\Delta S$  is \_\_\_\_\_ and  $\Delta H$  is \_\_\_\_\_.

- positive, negative
- negative, positive
- small, zero
- large, zero
- large, small

45. At what temperature would a given reaction become spontaneous if  $\Delta H = +119$  kJ and  $\Delta S = +263$  J/K?

- 452 K
- 2210 K
- 382 K
- 2.21 K
- 363 K

46. Given the following information, calculate  $\Delta G^\circ$  for the reaction below at 25 °C:



$\Delta H^\circ = 133.0$  kJ and  $\Delta S^\circ = 401.5$  J/K

- 252.6 kJ
- 13.4 kJ
- 13.4 kJ
- 122.9 kJ
- 252.6 kJ

47. For the process at 25 °C  $\text{I}_2(\text{g})$  to  $\text{I}_2(\text{s})$ , what are the signs of  $\Delta G$ ,  $\Delta H$ , and  $\Delta S$ ?


	$\Delta G$	$\Delta H$	$\Delta S$
a.	+	-	-
b.	-	-	-
c.	-	+	+
d.	-	-	+
e.	+	+	+

48. All of the following have  $\Delta G^\circ_f = 0$   
**EXCEPT**
- $O_2(g)$
  - $Br_2(g)$
  - $H_2(g)$
  - $Ca(s)$
  - $Hg(l)$
49. The disorder of a system is represented by the
- enthalpy.
  - Gibbs free energy.
  - entropy.
  - heat of vaporization.
  - equilibrium constant.
50. Calculate the standard entropy change for the following reaction,  
 $CH_4(g) + 2 O_2(g) \rightarrow CO_2(g) + 2H_2O(l)$   
 given that  $S^\circ[CO_2(g)] = 213.74$   
 $J/K \cdot mol$ ,  $S^\circ[O_2(g)] = 205.14$   
 $J/K \cdot mol$ ,  $S^\circ[H_2O(l)] = 69.91$   
 $J/K \cdot mol$ , and  $S^\circ[CH_4(g)] = 186.26$   
 $J/K \cdot mol$ .
- 312.89 J/K
  - 242.98 J/K
  - 118.42 J/K
  - 23.5 J/K
  - 312.89 J/K
51. Calculate the standard molar entropy of urea ( $CO(NH_2)_2(s)$ ) if the standard entropy change for the formation is  $-456.3 J/K \cdot mol$  and given  $S^\circ[C(s)] = 5.74 J/K \cdot mol$ ,  $S^\circ[O_2(g)] = 205.1 J/K \cdot mol$ ,  $S^\circ[N_2(g)] = 191.6 J/K \cdot mol$ , and  $S^\circ[H_2(g)] = 130.7 J/K \cdot mol$ .
- 1017.2 J/K•mol
  - +314.1 J/K•mol
  - +194.2 J/K•mol
  - +105.0 J/K•mol
  - 56.0 J/K•mol
52. For the reaction  
 $MgO(s) + CO_2(g) \rightarrow MgCO_3(s)$   
 $\Delta H^\circ_{rxn} = -178 \text{ kJ}$  and  $\Delta S^\circ_{rxn} = -161$   
 $J/mol \cdot K$ .
- Will the reaction be spontaneous at  $900^\circ C$ ?
- Yes, because  $\Delta G$  will change.
  - Yes, because  $\Delta H$  and  $\Delta S$  are temperature independent.
  - Yes, because  $\Delta H$  and  $\Delta S$  are temperature dependent.
  - No, because  $\Delta G$  is positive.
  - No, because  $\Delta G$  is negative.
53. If a process is exothermic and not spontaneous, then what must be true?
- $\Delta S > 0$
  - $\Delta H > 0$
  - $\Delta G = 0$
  - $\Delta S < 0$
  - $\Delta H = 0$
54. Which of the following is true about vaporization?
- $\Delta S$  is positive and  $\Delta H$  is negative.
  - $\Delta S$ ,  $\Delta H$ , and  $\Delta G$  are all negative.
  - $\Delta S$  and  $\Delta H$  are both negative.
  - $\Delta S$  and  $\Delta H$  are both positive.
  - $\Delta S$ ,  $\Delta H$ , and  $\Delta G$  are equal to zero.
55. Which of the following does not have a free energy of zero?
- $N_2(g)$
  - $I_2(s)$
  - $Fe(s)$
  - $Na(g)$
  - $He(g)$

56. Ammonium nitrate spontaneously dissolves in water at room temperature and the process causes the solution to become quite cold. Which of the following is **TRUE** about the dissolution of ammonium nitrate?

- a. The process is exothermic.
- b. Its solubility will be greater in warmer water.
- c.  $\Delta S^\circ$  for the reaction is negative.
- d. All solutions of ammonium nitrate are supersaturated.
- e. All solutions of ammonium nitrate are cold.

57. The following general reaction is not spontaneous at room temperature.



$$\Delta H^\circ = +50.0 \text{ kJ and } \Delta S^\circ = +100. \text{ J/K}$$

At what temperature will the reaction become spontaneous?

- a. 500 °C
- b. 0.5 K
- c. 500 K
- d. 250 °C
- e. Not at any temperature.